

Positive Invariance Lemmas for Control Problems with Convergence to Lyapunov-unstable Sets

A. Gorban,^{*} I. Tyukin,[†] E. Steur,[‡] and H. Nijmeijer[§]

January 22, 2009

Abstract

We provide Lyapunov-like characterizations of positive invariance, boundedness and convergence of non-trivial solutions for a class of systems with unstable invariant sets. The systems of this class comprise of a stable part coupled with a one-dimensional unstable or critically stable subsystem. Examples of these systems appear in the problems of nonlinear output regulation, parameter estimation and adaptive control. We demonstrate that, for a large class of systems with unstable equilibria and solutions that might escape to infinity in finite time, it is always possible to determine simple criteria for positive invariance and boundedness of the system's nontrivial solutions. Conversely, it is possible to characterize domains of initial conditions that lead to solutions escaping from the origin. In contrast to other works addressing convergence issues in unstable systems, our results do not rely on the availability of input-output gains or contraction rates that are usually required for the stable compartment.

Keywords: Convergence, weakly attracting sets, stability, Lyapunov functions, relaxation times, non-uniform small-gain theorem

1 Introduction

The analysis of asymptotic properties of dynamical systems is an important component of modern control theory. Even though the problems of control are often viewed as that of synthesis rather than analysis, the latter crucially affects the former. Indeed, in order to be able to specify feasible goals of synthesis, e.g. forward-completeness, state boundedness, asymptotic convergence of solutions to a region in the state space etc., one needs to understand how these properties depend on the system parameters and controls.

The majority of the analysis techniques in control, and hence methods for systems design, rely upon presumption that desired motions in the system are to stable in the

^{*}Dept. of Mathematics, University of Leicester, Leicester, LE1 7RH, UK (ag153@le.ac.uk)

[†]**Corresponding author.** Dept. of Mathematics, University of Leicester, Leicester, LE1 7RH, UK (I.Tyukin@le.ac.uk); Lab. for Perceptual Dynamics, RIKEN BSI, 2-1 Hirosawa, Wako-shi, Japan

[‡]Dept. of Mechanical Engineering, Eindhoven University of Technology, P.O. Box 513 5600 MB, Eindhoven, The Netherlands, (e.steur@tue.nl)

[§]Dept. of Mechanical Engineering, Eindhoven University of Technology, P.O. Box 513 5600 MB, Eindhoven, The Netherlands, (e-mail: h.nijmeijer@tue.nl)

sense of Lyapunov [7]. Let us briefly remind this and other related notions from the domain of dynamical systems.

Consider a flow $x(t, x_0) : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$. A set $\mathcal{S}$ is *invariant* w.r.t. $x(t, x_0)$ iff $x_0 \in \mathcal{S} \Rightarrow x(t, x_0) \in \mathcal{S}$ for all $t \in \mathbb{R}$. A set $\mathcal{S}$ is *positively (forward) invariant* iff $x(t_0, x_0) \in \mathcal{S} \Rightarrow x(t, x_0) \in \mathcal{S}$ for all $t \geq t_0$. Unions and intersections of a family of (positively) invariant sets are (positively) invariant. A closed invariant (w.r.t. the flow $x(t, x_0)$) set $\mathcal{S}$ is *stable in the sense of Lyapunov* if for any neighborhood $\mathcal{V}$ of $\mathcal{S}$ there exists a positively invariant (w.r.t the flow $x(t, x_0)$) neighborhood $\mathcal{W}$ of $\mathcal{S}$ such that $\mathcal{W} \subset \mathcal{V}$ [24]. In other words, a set that is stable in the sense of Lyapunov has a *fundamental base*¹ of positively invariant neighborhoods.

The notion of Lyapunov stability and analysis methods that are based on this notion are proven successful in a wide range of engineering applications including the problems of regulation, tracking, state observation [13], [5], identification [6] and adaptation [12], [10]. The popularity and success of the concept of Lyapunov stability resides, to a substantial degree, in the convenience and applied utility of the method of Lyapunov functions. Indeed, instead of explicit computation of solutions of an ordinary differential equation (ODE) it suffices to solve an algebraic inequality involving partial derivatives of a given Lyapunov candidate function.

Yet, as the methods of control expand from purely engineering applications into wider areas of science, there is a need for maintaining behavior that fail to obey the usual notion of uniform Lyapunov stability [19], [20]. Even though solutions of these systems may not be uniformly asymptotically stable (or even become unstable), they are required to be bounded.

In this paper we propose an extension of the classical Lyapunov function method that allows to address the issues of boundedness and convergence to invariant sets for a rich family of systems of which the solutions may not be uniformly asymptotically stable. Below we provide two examples of systems from this class that appear in the context of control and estimation problems.

Example 1 Nonlinear output regulation in presence of uncertainties. The studied question is as follows. Let the following system be given

$$\begin{aligned} \dot{x} &= f(x, t) + g(x, t)u, \quad f : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}^n, \quad g : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}^m \\ y &= h(x), \quad h : \mathbb{R}^n \rightarrow \mathbb{R}^d \end{aligned} \tag{1}$$

where x is a state vector and u is a function of which the range is in $\mathbb{R}^m$. What information about the system should be made available in order to be able to derive a feedback $u(h(x), t)$ that renders the solutions of (1) converge to zero asymptotically? The question has been answered in [4], [14]. In short, it was shown that 1) if $g(x, t)$ is bounded, 2) there exists a matrix $K \in \Omega_K \subset \mathbb{R}^{m \times d}$ such that the zero solution of

$$\dot{x} = f(x, t) + g(x, t)Kh(x)$$

is exponentially stable, and the bounds of Ω_K are known, then the feedback

$$\begin{aligned} u &= \beta(\gamma(\lambda))h(x), \\ \dot{\lambda} &= |h(x)|^p, \end{aligned} \tag{2}$$

¹A collection $\mathcal{U}_{\mathcal{S}}$ of all neighborhoods of $\mathcal{S}$ is called a *neighborhood system* of $\mathcal{S}$. An arbitrary subcollection $\mathcal{B}_{\mathcal{S}} \subset \mathcal{U}_{\mathcal{S}}$ is a *fundamental base* of system $\mathcal{U}_{\mathcal{S}}$ iff every neighborhood $V \in \mathcal{U}_{\mathcal{S}}$ contains at least one $B \in \mathcal{B}_{\mathcal{S}}$.

where $\beta : \mathbb{R} \rightarrow \mathbb{R}^{m \times d}$ is a dense trajectory in $\Omega_K \subset \mathbb{R}^{m \times d}$ and $\gamma : \mathbb{R} \rightarrow \mathbb{R}$ is a function that grows sufficiently slow, ensures that $\lim_{t \rightarrow \infty} x(t) = 0$. Notice that despite trajectories $x(t)$ converge to the origin asymptotically, solutions of the extended system (1), (2) are no longer uniformly asymptotically stable.

Example 2 Adaptive observers for systems in non-canonic adaptive observer form. Suppose that the following system be given

$$\begin{aligned} \dot{x}_1 &= \varphi_1(y, t)^T \theta_1 + \sum_{j=2}^n x_j + u(t), \\ \dot{x}_i &= -p_{i-1}x_i + \varphi_i(y, t)^T \theta_i, \quad i = 2, \dots, n, \quad p_i \in \mathbb{R}_{>0} \\ y &= (1, 0, \dots, 0)^T x, \end{aligned} \tag{3}$$

where $p = (p_1, \dots, p_{n-1}) \in \mathbb{R}^{n-1}$, $\theta_i = \text{col}(\theta_{i,1}, \dots, \theta_{i,m})$ are the vectors of unknown of unknown parameters, functions $\varphi_i : \mathbb{R}^d \times \mathbb{R} \rightarrow \mathbb{R}^m$, $u : \mathbb{R} \rightarrow \mathbb{R}^n$ are known functions. The problem is to infer the values of p_i , θ_i from the values $y(t)$. Standard Lyapunov-functions based techniques for estimating the values of p_i , θ_i , e.g. [9], do not apply here because system (3) is not in the canonical adaptive observer form (see [1], [8] for details). Yet, trading Lyapunov stability for convergence, allows to solve this problem. It has been shown in [23], [18] that there exists an estimator such that the dynamics of the extended system satisfies

$$\begin{aligned} \dot{z} &= A(t)z + g(z, p, t) - g(z, \beta(\lambda), t), \quad \beta : \mathbb{R} \rightarrow \mathbb{R}^{n-1} \\ \dot{\lambda} &= \gamma \|(1, 0, \dots, 0)^T z\|_\varepsilon, \quad \gamma \in \mathbb{R}_{>0}, \end{aligned} \tag{4}$$

with $z_1 = x_1$, $\dot{z} = A(t)z$ being uniformly asymptotically stable, $g(z, \beta(\lambda), t)$ being globally Lipschitz in λ respectively, and $\|s\|_\varepsilon = |s| - \varepsilon$ for $|s| \geq \varepsilon$ and $\|s\|_\varepsilon = 0$ for $|s| < \varepsilon$. In particular, $e(t) = z(t) - x(t)$ converges into a small neighborhood of the origin as $t \rightarrow \infty$. Furthermore, for any $\varepsilon > 0$ there exists an interval $(0, \gamma_0(\varepsilon)]$ such that $\limsup_{t \rightarrow \infty} \|\beta(\lambda(t)) - p\| \leq \delta(\varepsilon)$, $\delta \in \mathcal{K}$ for all $\gamma \in (0, \gamma_0]$.

It is easy to see that the invariant sets of (4) may not be uniformly asymptotically stable in the sense of Lyapunov. At the same time they are, as shown in [21], weakly attracting. As is mentioned in [11], control and estimation strategies similar to those presented in (1)–(4) can be viewed as a prototype for any pre-routing scheme in the domain of supervisory control. Hence understanding their limitations and capabilities is important for further progress in this field as well.

In all these examples asymptotically stable motion in a high-dimensional subspace is coupled with unstable explorative motion in a low-dimensional subspace. Even though solutions of these systems of ODE may not be stable in the sense of Lyapunov, they asymptotically converge to the required domains. Notwithstanding the relevance of such systems for modelling², control and identification, there are limitations restricting further progress in application of the broader concept of unstable convergence to these areas. Among these, from the authors point of view, is the lack of a simple analogue of the Lyapunov method for these, strictly speaking, unstable systems that would allow to draw

²Here we would like to refer to [16], [15] where dissipative saddles are used to model decision-making sequences.

conclusions about asymptotic properties of unstable solutions without the need of solving the equations.

Presently available techniques, such as e.g. non-uniform small-gain theorems [22], require existence of input-output gains for the stable subsystem. This condition, however, is already violated for simple cascade structures such as

$$\begin{aligned} \dot{x}_1 &= -kx_1 + x_1^2\lambda \\ \dot{\lambda} &= -\gamma x_1^2, \quad k, \gamma \in \mathbb{R}_{>0}, \end{aligned} \tag{5}$$

preventing the application of this theory. These techniques require knowledge, or at least estimates, of convergence rates for the stable component at $\lambda = 0$ [22]. Regarding the proofs of convergence in [4], they are of existential nature and also involve availability of the estimates (bounds) of the system solutions as functions of time. Moreover, as the authors of [4] comment, the results may not be of practical relevance in applications. Furthermore, they do not apply to system (5) for the term $x_1^2\lambda$ is not globally Lipschitz in x_1 . Hence, developing novel methods to address the issues of convergence to unstable sets is needed. These methods, on the one hand, should inherit the efficiency of Lyapunov analysis in which boundedness of trajectories can be verified by checking a system of inequalities without involving prior knowledge of the solutions of the system. On the other hand, these methods should apply to systems with instabilities. In our present contribution we provide a set of results that can be considered as a possible candidate.

We consider a class of systems which can be viewed as a generalization of (1)–(5). We chose this class of systems for its relevance in applications ranging from nonlinear regulation in control to the problems of modelling decision-making in neural systems. Since we are interested in the solutions that are not necessarily stable in the sense of Lyapunov we abandon the concept of neighborhoods from standard Lyapunov analysis [7], [24]. For a given invariant set $\mathcal{S}$ of a system, instead of searching for a fundamental base of positively invariant neighborhoods of $\mathcal{S}$, we study existence of an associated with $\mathcal{S}$ collection of positively invariant sets. These sets are not necessarily neighborhoods, and they are not required to form a fundamental base. In particular, the sets are allowed to be closed, and their boundaries may have non-empty intersections with $\mathcal{S}$.

For the chosen class of dynamical systems we formulate Lyapunov-like conditions that allow to specify the domains of positive invariance containing Lyapunov-unstable equilibria on their boundaries. In the classical method of Lyapunov functions the role of a Lyapunov function is to assure that an invariant set, e.g. an equilibrium, has a fundamental base of positively invariant neighborhoods. In our work we use an extension of this method in which a substitute of a Lyapunov function is used to demonstrate existence of a family of positively invariant sets (not necessarily neighborhoods) associated with the equilibrium. These results allow us to estimate the domains of initial conditions, as functions of system parameters in problems (1)–(5), that lead to bounded solutions without the need to know convergence rate of the stable part of (1)–(5). Parameters of these systems are not required to be known precisely, and input-output gains of the systems need not to be defined. Furthermore, in contrast to our previous results on the same topic [22], present conditions allow to specify domains of initial conditions that lead to trajectories necessarily escaping a neighborhood of the equilibria in question.

The paper is organized as follows. In Section 2 we formulate the problem and specify main assumptions in detail. In Section 3 we provide main results of the paper, and Section 4 contains brief discussion and conclusions. The following notational conventions are used

throughout the paper: symbol $\mathcal{C}^k$ denotes the space of functions that are at least k times differentiable; $\|\cdot\|$ stands for the Euclidian norm, $\mathcal{P}$ denotes the class of positive-valued continuous functions $p : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ such that $p(0) = 0$ and $p(s) > 0$ for all $s > 0$; $\mathcal{K}$ denotes the class of all strictly increasing continuous functions $\kappa : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ such that $\kappa(0) = 0$. If, in addition, $\lim_{s \rightarrow \infty} \kappa(s) = \infty$ we say that $\kappa \in \mathcal{K}_{\infty}$.

2 Problem Formulation

Consider the following system:

$$\begin{aligned}\dot{x} &= f(x, \lambda, t) \\ \dot{\lambda} &= g(x, \lambda, t),\end{aligned}\tag{6}$$

where the functions

$$\begin{aligned}f &: \mathbb{R}^n \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}^n, \\ g &: \mathbb{R}^n \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}\end{aligned}$$

are continuous and satisfy Assumptions 1–2 below.

Assumption 1 *The function $f(\cdot, \cdot, \cdot)$ in (6) is locally Lipschitz in x and λ uniformly in t , and there exists $V : \mathbb{R}^n \rightarrow \mathbb{R}$, $V \in \mathcal{C}^1$ such that*

$$\begin{aligned}\underline{\alpha}(\|x\|) \leq V(x) \leq \bar{\alpha}(\|x\|), \quad \underline{\alpha}, \bar{\alpha} \in \mathcal{K}_{\infty} \\ \frac{\partial V}{\partial x} f(x, \lambda, t) \leq \alpha(V(x)) + \beta(V(x))\varphi(|\lambda|), \quad \alpha, \beta \in \mathcal{C}^0, \quad \varphi \in \mathcal{K}.\end{aligned}\tag{7}$$

Assumption 1 holds, for example, for those systems in which the term $(\partial V / \partial x) f(x, \lambda, t)$ can be bounded from above as follows:

$$\frac{\partial V}{\partial x} f(x, \lambda, t) \leq -\alpha_0(\|x\|) + \beta_0(\|x\|)\varphi(|\lambda|), \quad \alpha_0, \beta_0 \in \mathcal{K}.\tag{8}$$

Indeed, (7) follows immediately from (8) with $\alpha(\cdot) = -\alpha_0(\bar{\alpha}^{-1}(\cdot))$, $\beta(\cdot) = \beta_0(\underline{\alpha}^{-1}(\cdot))$, and $\underline{\alpha}^{-1}(\cdot)$, $\bar{\alpha}^{-1}(\cdot)$ are the inverses of $\underline{\alpha}(\cdot)$, $\bar{\alpha}(\cdot)$ respectively. Notice also that in this case Assumption 1 states that the origin of $\dot{x} = f(x, 0, t)$ is globally stable in the sense of Lyapunov, and V is the corresponding Lyapunov function. Even though the right-hand side of (6) is allowed to be time-varying, we restrict our consideration to systems for which the function V does not depend on time explicitly.

Assumption 2 *The function $g(\cdot, \cdot, \cdot)$ in (6) is locally Lipschitz in x and λ uniformly in t , and there exist $\delta, \xi \in \mathcal{K}$ such that the following inequality holds for all $x \in \mathbb{R}^n$, $t \in \mathbb{R}$:*

$$-\xi(|\lambda|) - \delta(\|x\|) \leq g(x, \lambda, t) \leq 0 \quad \forall \lambda \geq 0.\tag{9}$$

Assumption 2 reflects the fact that derivative $\dot{\lambda}$ does not change sign for all $\lambda \geq 0$. Without loss of generality we consider the case when λ is non-increasing with time, and alternative formulations of our conclusions are available if (9) is replaced with

$$0 \leq g(x, \lambda, t) \leq \delta(\|x\|) + \xi(|\lambda|) \quad \forall \lambda \leq 0.\tag{10}$$

We aim to formulate a set of conditions that would allow us to estimate domains of positive invariance of (6) and specifically those in which the solutions of (6) remain bounded. These conditions are provided in the next section.




Figure 1: Illustration of the proof of Lemma 1. Panel *a*: boundary $\lambda = \psi(V)$. The vector $(\partial\psi/\partial V, -1)$ is a normal vector to the curve $\lambda = \psi(V)$ at the point *A*. Because $\partial\psi/\partial V \geq 0$ it is always pointing in the direction of $\lambda < \psi(V)$. Panel *b*: boundary $\lambda = \text{const}$. Because $\dot{\lambda} \leq 0$ for all $\lambda \geq 0$ the vector $(dV, d\lambda)$ is pointing in the direction of $\lambda \geq \psi(V)$.

3 Main Results

We start with the following lemma

Lemma 1 (Boundedness 1) *Let system (6) be given and satisfy Assumptions 1, 2. Suppose that there exist a function*

$$\psi : \psi \in \mathcal{K} \cap \mathcal{C}^1$$

and a positive constant $a \in \mathbb{R}_{>0}$ such that

$$\frac{\partial\psi(V)}{\partial V} [\alpha(V) + \beta(V)\varphi(\psi(V))] + \delta(\underline{\alpha}^{-1}(V)) + \xi(\psi(V)) \leq 0, \quad \forall V \in [0, a]. \quad (11)$$

Then the domain

$$\Omega_a = \{(x, \lambda) \mid x \in \mathbb{R}^n, \lambda \in \mathbb{R}_{\geq 0}, \psi(a) \geq \lambda \geq \psi(V(x)), V(x) \in [0, a]\} \quad (12)$$

is forward invariant with respect to (6), and furthermore

$$\exists \lambda' \in [0, \psi(a)] : \lim_{t \rightarrow \infty} \lambda(t) = \lambda', \quad (13)$$

$$\lim_{t \rightarrow \infty} \|x(t)\| = 0. \quad (14)$$

Proof of Lemma 1. The main idea behind the proof is as follows (see also Fig. 1). Consider the surface

$$\lambda = \psi(V(x)).$$

The surface splits the system's state space into two domains: $\lambda < \psi(V(x))$ (white area in Fig. 1) and $\lambda \geq \psi(V(x))$ (gray area in Fig. 1). Let us pick a non-empty interval $[0, a]$, $a \in \mathbb{R}_{>0}$ and set $0 \leq \lambda(t_0) \leq \psi(a)$. Then $V(x(t))$ and $\lambda(t)$ remain in Ω_a specified by (12) (gray area in Fig. 1) if the right-hand side of

$$\begin{aligned} \dot{V} &= \frac{\partial V}{\partial x} f(x, \lambda, t) \\ \dot{\lambda} &= g(x, \lambda, t) \end{aligned}$$

is pointing inside the domain $\lambda \geq \psi(V(x))$ for all $V \in [0, a]$, $\lambda = \psi(V)$. According to the assumptions, function $\psi \in \mathcal{K} \cap \mathcal{C}^1$, i.e. it is increasing and continuously differentiable. Hence the vector $(\partial\psi/\partial V, -1)$ is (locally) orthogonal to the curve $\lambda = \psi(V)$ for all $V \geq 0$. Therefore the fact that the dot product $(\partial\psi/\partial V, -1)(dV, d\lambda)^T$ is non-positive at $\lambda = \psi(V)$, $V \in [0, a]$, or alternatively, that the following condition holds:

$$\left. \frac{\partial\psi}{\partial V} \dot{V} - \dot{\lambda} \right|_{\lambda=\psi(V)} \leq 0, \quad \forall V \in [0, a], \quad (15)$$

suffices to guarantee positive invariance of Ω_a . The latter property together with (7) automatically implies that $\|x(t, x_0, \lambda_0)\| \leq \underline{\alpha}^{-1}(a)$, $|\lambda(t, x_0, \lambda_0)| \leq \psi(a)$ for all $(x_0, \lambda_0) \in \Omega_a$.

According to Assumptions 1, 2, taking into account that $\psi, \varphi, \delta, \xi$ are non-decreasing, and noticing that $\lambda = \psi(V(x)) \geq 0$ for all $V(x) \in [0, a]$ we have:

$$\begin{aligned} \left. \frac{\partial\psi}{\partial V} \dot{V} - \dot{\lambda} \right|_{\lambda=\psi(V(x))} &= \left. \frac{\partial\psi}{\partial V} \frac{\partial V}{\partial x} f(x, \lambda, t) - g(x, \lambda, t) \right|_{\lambda=\psi(V(x))} \\ &\leq \left. \frac{\partial\psi}{\partial V} [\alpha(V(x)) + \beta(V(x))\varphi(|\lambda|)] + \delta(\|x\|) + \xi(|\lambda|) \right|_{\lambda=\psi(V(x))} \\ &\leq \frac{\partial\psi}{\partial V} [\alpha(V(x)) + \beta(V(x))\varphi(\psi(V(x)))] + \delta(\underline{\alpha}^{-1}(V(x))) + \xi(\psi(V(x))) \end{aligned} \quad (16)$$

Hence (11) implies that (15) holds, and boundedness of $x(t, x_0, \lambda_0)$, $\lambda(t, x_0, \lambda_0)$ for all $(x_0, \lambda_0) \in \Omega_a$ follows.

Property (13) is immediate consequence of Bolzano-Weierstrass theorem, and (14) follows from Barbalat lemma. $\square$

Remark 1 The conditions of Lemma 1 imply that $\psi(V)$ is non-increasing with time. Hence $d\psi(V)/dt \leq 0$ for all $V \in [0, a]$ may serve as a necessary condition for the positive invariance of Ω_a .

Example 3 Let us illustrate the lemma with deriving the domains of positive invariance for (5). As we mentioned earlier, the first equation in (5) does not have finite input-output gain with respect to the input-output pair (x_1, λ) . Hence neither of the approaches presented in [4], [22] apply. We shall show now that Lemma 1 allows to determine domains of initial conditions for (5) corresponding to bounded solutions without invoking a finite input-output gain requirement.

In order to apply Lemma 1 we need to check Assumptions 1 and 2. Assumption 1 holds for (5) with $V = x_1^2$, $\alpha(V) = -k2V$, $\beta(V) = 2V^{3/2}$, $\varphi(|\lambda|) = |\lambda|$, and Assumption 2 holds with $\delta(\|x\|) = \gamma x_1^2$ and $\xi = 0$. Now we need to choose a candidate for $\psi(V)$. For simplicity, let

$$\psi(V) = pV, \quad p \in \mathbb{R}_{>0}.$$

With this function condition (11) reduces to the following constraint

$$-2kp + 2p^2V^{3/2} + \gamma \leq 0 \quad \forall V \in [0, a]$$

This constraint is satisfied for all $V \in [0, a]$ with a defined as:

$$a = \left(\frac{1}{p} \left(k - \frac{\gamma}{2p} \right) \right)^{2/3}, \quad p \geq \frac{\gamma}{2k} > 0$$




Figure 2: Domain of positive invariance for (5). Gray area shows the union of domains $\Omega_a(p)$ for all $p \geq \frac{\gamma}{2k}$. Examples of elementary $\Omega_a(p_i)$, $p_1 = 0.3$, $p_2 = 3$, $p_3 = 20$ are depicted as shaded areas.

Hence domains of positive invariance for (5) shall contain the following union of cones

$$\cup_{p \geq \frac{\gamma}{2k}} \left\{ \Omega_a(p) = \{(x_1, \lambda) \mid \lambda \geq p|x_1|^2, 0 \leq |x_1| \leq \left(\frac{1}{p} \left(k - \frac{\gamma}{2p} \right) \right)^{1/3} \} \right\}$$

For $\gamma = 0.1$ and $k = 1$ this union is depicted in Fig. 2 $\square$

Remark 2 Lemma 1 requires that function $\psi(V)$ is strictly increasing in V and differentiable. In principle, one can remove this restriction by considering $\psi(V)$ as a piece-wise monotone function and replacing (11) with a set of similar inequalities for each segment of monotonicity. The only difference would be that the sign of these inequalities will have to vary depending on the type of monotonicity of $\psi(V)$ in the segment (i.e. increasing/decreasing).

As for the requirement of differentiability, it can be removed if we use a slightly different argument for establishing existence of the domains of positive invariance. The argument involves a notion of a star shaped domain. Formal definition of this and other related notions, and basic properties of star shaped domains and functions are provided in Appendix.

Let us remind that a domain $\Omega \subset \mathbb{R}^n$ is called *star-shaped with respect to the origin* if every segment connecting the origin with $x \in \Omega$ lies entirely in Ω . Clearly star shapedness of a set is a weaker property than convexity, and our statements involving the notion of star shapedness should hold if “star shaped” is replaced with “convex” in their formulations. A star shaped envelop of a set D (w.r.t. the origin) is a minimal star shaped set (w.r.t. the origin) including D . This set is denoted as:

$$\text{star}_x(D).$$

The *epigraph* of a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$, or simply $\text{epi}(f)$, is the domain in $\mathbb{R}^{n+1}$: $\text{epi}(f) = \{x \in \mathbb{R}^n, \mu \in \mathbb{R} \mid f(x) \leq \mu\}$. A function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is called *star shaped* with respect to the origin iff its epigraph is a star shaped domain with respect to the origin. Convex




Figure 3: Deriving conditions of positive invariance for the domains bounded by non-differentiable functions $\lambda = \psi(V)$. In this case the fact that the epigraph $\text{epi}(\psi)$ of the function $\psi(V)$ over $[0, a]$ is star-shaped with respect to the origin and that $\tau(t)$ is non-increasing ensure that trajectories $V(x(t)), \lambda(t)$ will not pass through the boundary $\lambda = \psi(V)$. Star-shapedness of the $\text{epi}(\psi)$ is needed to guarantee that the vector $(V(t_1) - V(t_2), \lambda(t_1) - \lambda(t_2))$ points in the direction of $\psi(V) < \lambda$ for all points on the curve $\lambda = \psi(V), V \in [0, a]$. Notice that this condition does not hold for the point B .

envelope of a function f is denoted as $\text{conv}(f)(x)$ and star shaped envelop of f (w.r.t. the origin) is denoted as $\text{star}(f)(x)$. In addition we define

$$\text{epi}(f_{[0,a]}) = \{(x, \mu) \mid x \in [0, a]^n, \mu \in \mathbb{R}, f(x) \leq \mu\},$$

the epigraph of the restriction of f on the hypercube $[0, a]^n$, and consider convex and star-shaped envelops of such restriction:

$$\text{conv}(f_{[0,a]})(x) = \inf\{\mu \mid (x, \mu) \in \text{conv epi}(f_{[0,a]})\}$$

$$\text{star}(f_{[0,a]})(x) = \inf\{\mu \mid (x, \mu) \in \text{star epi}(f_{[0,a]})\}.$$

In order to derive domains of positive invariance of which the boundaries are not necessarily differentiable curves we use the following strategy. Instead of searching for the domains in which the dot product $(\partial\psi/\partial V, -1)(dV, d\lambda)^T$ is non-positive we search for an interval $[0, a]$ such that the epigraph of ψ for $V \in [0, a]$:

$$\text{epi}(\psi_{[0,a]}) = \{(V, \lambda) \mid V \in [0, a], \lambda \in \mathbb{R}_{\geq 0}, \psi(V) < \lambda\}$$

is star-shaped with respect to the origin, and ratio $V(x)/\lambda$ is not-increasing in forward time for all $V \in [0, a], \lambda = \psi(V)$ (see also Fig. 3). In other words

$$\left. \frac{d}{dt} \left(\frac{V}{\lambda} \right) \right|_{\lambda=\psi(V)} \leq 0, \forall V \in [0, a]. \quad (17)$$

Inequality (17) implies that the curve $\lambda = \psi(V)$ "shrinks" locally in the direction of $V = 0$ for every pair $(V, \lambda), \lambda = \psi(V), V \in [0, a]$. Hence trajectories $V(x(t)), \lambda(t)$ passing through Ω_a at $t = t_0$ will remain there for all $t \geq t_0$. Therefore the following statement holds.

Lemma 2 (Boundedness 2) *Let system (6) be given and satisfy Assumptions 1, 2. Suppose that there exist a function $\psi : \mathbb{R} \rightarrow \mathbb{R}$, $\psi(0) = 0$, and a positive constant $a \in \mathbb{R}_{>0}$ such that*

$$\psi(V) [\alpha(V) + \beta(V)\varphi(\psi(V))] + V [\delta(\underline{\alpha}^{-1}(V)) + \xi(\psi(V))] \leq 0, \quad \forall V \in [0, a], \quad (18)$$

and $\text{epi}(\psi|_{[0,a]})$ is star-shaped with respect to the origin.

Then domain (12) is forward invariant with respect to (6), and furthermore properties (13), (14) hold.

Proof of Lemma 2. The proof is straightforward and follows from (17). Inequality (17) is satisfied if

$$\dot{V}\lambda - \dot{\lambda}V \Big|_{\lambda=\psi(V)} = \psi(V)\dot{V} - \dot{\lambda}V \leq 0$$

On the other hand, similar to (16), the following estimate holds:

$$\begin{aligned} \psi(V)\dot{V} - \dot{\lambda}V &\leq \\ \psi(V) [\alpha(V(x)) + \beta(V(x))\varphi(\psi(V(x)))] + V [\delta(\underline{\alpha}^{-1}(V(x))) + \xi(\psi(V(x)))] &= \mathcal{G}(V(x)) \end{aligned}$$

Therefore, $\mathcal{G}(V(x)) \leq 0 \Rightarrow d/dt(V/\lambda) \leq 0$ at $\lambda = \psi(V)$. The remaining part of the proof is identical to that of Lemma 1. $\square$

Remark 3 The major difference between Lemmas 1 and 2 is that the partial derivative $\partial\psi/\partial V$ in (11) is replaced with the ratio $\psi(V)/V$ in the conditions that follow from (18). In case of linear functions $\psi = pV$ these lemmas will produce identical estimates of the domains of positive invariance. In the nonlinear case Lemma 1 allows to take the local structure of $\psi(V)$ into account while Lemma 2 does not. The price for this benefit is the required differentiability of $\psi(V)$.

Remark 4 Lemmas 1 and 2 establish criteria for existence of positively invariant sets in the state space of (6) in terms of the properties of functions $f(x, \lambda, t)$ and $g(x, \lambda, t)$. In control problems the function $f(x, \lambda, t)$ defines a controlled dynamical system, and $g(x, \lambda, t)$ describes tuning mechanisms of the controller (see, for instance Examples 1, 2). It is therefore important to know if, for a given controlled system

$$\dot{x} = f(x, \lambda, t),$$

where λ is the controller parameter, one can determine a tuning algorithm

$$\dot{\lambda} = g(x, \lambda, t) \quad (19)$$

ensuring existence of the domains of positive invariance corresponding to the bounded solutions in forward time. Results are available for the functions $f(x, \lambda, t)$ that are globally Lipschitz in λ uniformly in t [4], [22]. Lemma 2 allows to demonstrate that it is possible to derive tuning algorithms (19) ensuring existence of the domains of initial conditions that correspond to bounded solutions of (6) in which the right-hand side is only *locally* Lipschitz in λ . In particular, the following Corollary holds.

Corollary 1 (Existence) *Let the following system be given,*

$$\begin{aligned}\dot{x} &= f(x, \lambda, t) \\ \dot{\lambda} &= u,\end{aligned}\tag{20}$$

where u is a control input, and the function $f : \mathbb{R}^n \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}^n$ satisfies Assumption 1 with $\varphi(\cdot)$ being locally Lipschitz, and $\alpha(\cdot) = -\alpha_0(\cdot)$, $\alpha_0 \in \mathcal{P}$.

Then there exists a continuous function $g(x, \lambda, t) : \mathbb{R}^n \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ and a domain Ω_a with non-zero volume³ such that all solutions of (20) passing through Ω_a are bounded in forward time and, in addition, satisfy (13), (14).

Proof of Corollary 1. Let $\mathcal{S}(a)$ denote a class of functions from $\mathcal{P}$ such that their restriction to $[0, a]$ is star-shaped. According to Lemma 2, existence of functions $\psi(V) \in \mathcal{S}(a)$, $\gamma(V) \in \mathcal{P}$ such that the following inequality holds

$$\psi(V) [-\alpha_0(V) + \beta(V)\varphi(\psi(V))] + V\gamma(V) \leq 0, \quad \forall V \in [0, a], \quad a \in \mathbb{R}_{>0} \tag{21}$$

constitutes the proof. Let us show that for any function $f(x, \lambda, t)$ in (20) satisfying the assumptions of the corollary there will always exist functions $\psi(V) \in \mathcal{S}(a)$, $\gamma(V) \in \mathcal{P}$ and $a \in \mathbb{R}_{>0}$ such that inequality (21) holds.

First, we notice that a product of two functions from $\mathcal{P}$ belongs to $\mathcal{P}$. Hence we can always choose the function $\gamma(V)$ in the following form $\gamma(V) = \psi(V)\gamma_0(V)$, where $\gamma_0(\cdot)$ is some function from $\mathcal{P}$. Taking into account that the function $\varphi(V)$ is locally Lipschitz and that $\beta(V)$ is continuous we can observe that for any $a \in \mathbb{R}_{>0}$ there exists a positive constant $D(a) \in \mathbb{R}_{>0}$ such that

$$\beta(V)\varphi(\psi(V)) \leq D(a)\psi(V) \quad \forall V \in [0, a].$$

Therefore all we need to show is that for some $a \in \mathbb{R}_{>0}$ there exist $\psi(V) \in \mathcal{S}(a)$, $\gamma_0(V) \in \mathcal{P}$ such that the following inequality holds:

$$-\alpha_0(V) + D(a)\psi(V) + a\gamma_0(V) \leq 0, \quad \forall V \in [0, a]. \tag{22}$$

Choosing

$$\gamma_0(V) = \frac{\alpha_0(V)}{2a}$$

and taking (22) into account we obtain the following inequality

$$-\frac{1}{2}\alpha_0(V) + D(a)\psi(V) \leq 0, \quad \forall V \in [0, a]. \tag{23}$$

Therefore the corollary will be proven if we demonstrate that for every $\eta(V) \in \mathcal{P}$ there exists a star-shaped $\psi(V) \in \mathcal{P}$ and $a \in \mathbb{R}_{>0}$ such that

$$\eta(V) \geq \psi(V) \quad \forall V \in [0, a].$$

Existence of such function $\psi(V)$ is guaranteed by Proposition 1.

³In fact we show that Ω_a can be determined as in (12) where $\psi(\cdot)$ is a function from $\mathcal{K}_\infty$ and a is a positive number.

According to Proposition 1, for every $\eta(V) \in \mathcal{P}$ there exists $\text{star}(\eta_{[0,a]})(x)$. The function $\text{star}(\eta_{[0,a]})(x)$ is a continuous and monotone function, and it is a minorant of $\eta(x)$ on $[0, a]$. Hence choosing either

$$\psi(V) = \text{star} \left(\frac{1}{2D(a)} \alpha_{0 \ [0,a]} \right) (V)$$

or

$$\psi(V) = \text{conv} \left(\frac{1}{2D(a)} \alpha_{0 \ [0,a]} \right) (V)$$

we ensure that (23) holds with a star-shaped $\psi(V)$ from $\mathcal{P}$. Thus the corollary follows immediately from Lemma 2. $\square$.

Let us illustrate the utility of Corollary 1 with the following example.

Example 4 Consider the following nonlinear local regulation problem in presence of parametric uncertainty. Suppose that the system equations are defined as

$$\dot{x} = -\kappa(x) + (x^2 + 1)f(x, \theta) + u, \quad (24)$$

where

$$\kappa(x) = \begin{cases} 1, & x > 1 \\ x^3, & x \in [-1, 1] \\ -1, & x < -1 \end{cases}$$

$f(x, \theta)$ is a Lipschitz function with respect to θ :

$$|f(x, \theta) - f(x, \theta')| \leq D|\theta - \theta'|,$$

and θ is an unknown parameter from $\mathbb{R}$. The question is whether for some interval $\Omega_\theta \in \mathbb{R}$ there exists a compensating feedback

$$\begin{aligned} u(x, \hat{\theta}) &= -(x^2 + 1)f(x, \hat{\theta}) \\ \dot{\hat{\theta}} &= g(x), \quad g \in \mathcal{C}^1 \end{aligned}$$

and a domain of initial conditions, Ω_x , with positive measure such that trajectories $x(t)$ of (24) passing through Ω_x converge to the origin for all $\theta \in \Omega_\theta$.

Notice that because the zero solution of $\dot{x} = -\kappa(x)$ is not exponentially stable we cannot apply conventional theorems from the domain of output regulation [2]. Yet, as follows from Corollary 1 such feedback indeed exists. In order to see this consider the following functions $V(x) = 1/2x^2$, $\lambda = \theta - \hat{\theta}$. Clearly, Assumption 1 is satisfied for (24) with

$$\alpha_0(V) = \kappa(\sqrt{V}2)\sqrt{V}2, \quad \beta(V) = 2D\sqrt{V}(2V + 1),$$

and $\alpha_0(V), \beta(V) \in \mathcal{P}$. Hence for any θ there exists a domain $\Omega_a(\theta)$ with nonzero volume such that solutions passing through $\Omega_a(\theta)$ have the desired property: $\lim_{t \rightarrow \infty} x(t) = 0$. The intersection of $\Omega_a(\theta)$, $\theta \in \Omega_\theta$ is a non-empty set with nonzero volume provided that Ω_θ is sufficiently small.

The control scheme in example 4 can be generalized further to constitute an existence result to the problem of local nonlinear adaptive regulation. This simple generalization is provided in the next corollary.

Corollary 2 (Local adaptive regulation) *Consider the following system*

$$\dot{x} = f(x, \theta, t) + v, \quad (25)$$

where $f : \mathbb{R}^n \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}^n$ is a continuous function that is Lipschitz in θ locally in x

$$\|f(x, \theta, t) - f(x, \theta', t)\| \leq D(\|x\|)|\theta - \theta'|, \quad D \in \mathcal{K}$$

and v is a control input defined as

$$v = -f(x, \hat{\theta}, t) + f_0(x).$$

Suppose that $f_0(x)$ is a continuous vector-field such that the origin of $\dot{x} = f_0(x)$ is globally asymptotically stable in the sense of Lyapunov.

Then there exists a tuning algorithm:

$$\dot{\hat{\theta}} = g(x), \quad g(x) \in \mathcal{C}^0 \quad (26)$$

and a set of initial conditions Ω_a with non-zero volume such that for all $\theta \in [a, b]$

1) solutions of (25), (26) passing through Ω_a are bounded

2) $x(t)$ is asymptotically stirred to the origin: $\lim_{t \rightarrow \infty} \|x(t)\| = 0$

provided that $[a, b]$ is sufficiently small.

Remark 5 Solutions to the local adaptive regulation problem have been already discussed in the literature, e.g. in the framework of robust output regulation [2]. These results, however, require that the origin of $\dot{x} = f_0(x)$ is exponentially stable. Our present conclusions demonstrate that for a general class of systems, existence of an asymptotically stabilizing feedback suffices to show existence of an adaptive controller solving the problem of regulation locally in the presence of locally Lipschitz nonlinearly parameterized uncertainty.

Next, we characterize the domains of initial conditions corresponding to solutions that do not converge to the origin. These domains are provided in the following lemma.

Lemma 3 (Solutions escaping the origin) *Let system (6) be given and satisfy Assumptions 1, 2. Suppose that there exists a function*

$$\psi : \psi \in \mathcal{K} \cap \mathcal{C}^1$$

and a positive constant $a \in \mathbb{R}_{>0}$ such that for some $\varepsilon \in \mathbb{R}_{>0}$ the following holds

$$\frac{\partial \psi(V)}{\partial V} [\alpha(V) + \beta(V)\varphi(\psi(V) - \varepsilon)] + \delta(\underline{\alpha}^{-1}(V)) + \xi(\psi(V) - \varepsilon) \geq 0 \quad \forall V \in [0, a] \quad (27)$$

Then the domain

$$\Omega_a = \{(x, \lambda) \mid x \in \mathbb{R}^n, \lambda \in \mathbb{R}, \lambda \leq \psi(V(x)) - \varepsilon, V(x) \in [0, a]\} \quad (28)$$

is forward invariant with respect to (6). In other words, solutions of (6) passing through Ω_a do not converge to the origin.




Figure 4: Illustration to the proof of Lemma 3. Panel *a*: boundary $\lambda = \psi(V) - \varepsilon$. Vector $(\partial\psi/\partial V, -1)$ is a normal vector to the curve $\lambda = \psi(V) - \varepsilon$ at the point *A*. Because $\partial\psi/\partial V \geq 0$ it is always pointing into the direction of $\lambda < \psi(V)$. Panel *b*: boundary $\lambda = \text{const}$. Because $\dot{\lambda} \leq 0$ for all $\lambda \geq 0$ vector $(dV, d\lambda)$ is pointing into the direction of $\lambda \geq \psi(V)$.

Proof of Lemma 3 is analogous to that of Lemma 1 except that constraint $\lambda = \psi(V(x))$ is replaced with

$$\lambda = \psi(V(x)) - \varepsilon,$$

and the sign of (15) should be reversed (see also Fig. 4). $\square$

Example 5 Let us illustrate the lemma with system (5). Choosing the same functions V and $\psi(V)$ as in Example 3 we can rewrite inequality (27) as follows:

$$-2kp + 2pV^{1/2}(pV - \varepsilon) + \gamma \geq 0 \quad \forall V \in [0, a] \quad (29)$$

The term $2pV^{1/2}(pV - \varepsilon)$ is bounded from below by $-4/3\sqrt{\varepsilon^3 p/3}$. Hence condition (29) will be satisfied for all $V \geq 0$ if the values of p are chosen to obey the following inequality

$$-2kp - 4/3\sqrt{\varepsilon^3/3}\sqrt{p} + \gamma \geq 0.$$

Thus solutions passing through the following domain

$$\lambda \leq p|x_1|^2 - \varepsilon, \quad p \in \left(0, \left[\frac{\sqrt{\frac{16}{27}\varepsilon^3 + 8\gamma k} - \frac{4}{3}\sqrt{\frac{\varepsilon^3}{3}}}{4k}\right]^2\right)$$

will not converge to the origin.

So far we have established conditions that allow to derive Lyapunov-like characterizations of the domains of positive invariance for (6). These conditions do not require a-priori knowledge of the system solutions, nor they require estimation of an input-output gain of the stable part of the cascaded system. As we have seen, e.g. in the case of Lemma 1, the final shape of the domain of positive invariance can be estimated as a union of parameterized families of cones (see Fig. 2). Furthermore, these estimates can be obtained even for systems with finite escape time.

An important question remains: how good are these estimates in comparison with that of [22], [4] derived for the same problems. In order to answer this question consider the following subclass of (6) to which the results of [22], [4] apply:

$$\begin{aligned}\dot{x} &= f(x, t) + b(x, \lambda, t) \\ \dot{\lambda} &= -\gamma\|x\|, \quad \gamma \in \mathbb{R}_{>0}\end{aligned}\tag{30}$$

Functions $f : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}^n$, $b : \mathbb{R}^n \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}^n$, $g : \mathbb{R}^n \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}^n$ in (30) are assumed to be continuous; the function $b(x, \lambda, t)$ is Lipschitz in λ :

$$\|b(x, \lambda, t) - b(x, \lambda', t)\| \leq B|\lambda - \lambda'| \tag{31}$$

with $b(x, 0, t) = 0$. For simplicity, suppose that there exists a quadratic Lyapunov function

$$V(x) = x^T P x, \quad P = P^T > 0, \quad \underline{p}\|x\|^2 \leq V(x) \leq \bar{p}\|x\|^2, \quad \|P x\| \leq \sigma\|x\|, \quad \bar{p}, \underline{p} \in \mathbb{R}_{>0}$$

such that

$$\frac{\partial V(x)}{\partial x} f(x, t) \leq -kV, \quad k \in \mathbb{R}_{>0}. \tag{32}$$

We will assume that the values of $\bar{p}$, $\underline{p}$, k , and σ (maximal eigenvalue σ of P) are known. Then domains of initial conditions corresponding to bounded solutions of (30) are specified in the following corollary from Lemma 1

Corollary 3 (Non-uniform small-gain condition) *Let system (30) be given and assumptions (31), (32) hold. Then solutions of (30) passing through*

$$\Omega'_a = \{(x, \lambda) \mid x \in \mathbb{R}^n, \lambda \in \mathbb{R}, \lambda(t_0) \geq \|x(t_0)\|k\sqrt{\underline{p}\bar{p}}/(4\sigma B)\}$$

are bounded and remain in

$$\Omega_a = \{(x, \lambda) \mid x \in \mathbb{R}^n, \lambda \in \mathbb{R}, \lambda(t_0) \geq \|x(t_0)\|k\underline{p}/(4\sigma B)\} \supset \Omega'_a$$

provided that γ is sufficiently small:

$$\gamma < \frac{k^2}{16} \left(\frac{\underline{p}}{\sigma B} \right). \tag{33}$$

Proof of Corollary 3. In order to apply Lemma 1 we need to check Assumption 1 and 2. The first inequality in Assumption 1 clearly holds with

$$\bar{\alpha}(\|x\|) = \bar{p}\|x\|^2, \quad \underline{\alpha}(\|x\|) = \underline{p}\|x\|^2.$$

In order to check that the second inequality holds consider

$$\frac{\partial V}{\partial x}(f(x, t) + b(x, \lambda, t)) \leq -kV + 2\|P x\|\|b(x, \lambda, t)\| \leq -kV + \left(\frac{2\sigma B}{\sqrt{\underline{p}}} \right) \sqrt{V}|\lambda| \tag{34}$$

Hence the second inequality of Assumption 1 holds with

$$\alpha(V) = -kV, \quad \beta(V) = c\sqrt{V}, \quad c = \left(\frac{2\sigma B}{\sqrt{\underline{p}}} \right), \quad \varphi(|\lambda|) = |\lambda|.$$

Assumption 2 holds with $\delta(\|x\|) = \gamma\|x\|$ and $\xi(\|x\|) = 0$. According to Lemma 1, we need to find $\psi(V)$ such that (11) holds. Then the domains of positive invariance and boundedness of solutions are determined by (12). Let us choose

$$\psi(V) = \sqrt{V}r, \quad r \in \mathbb{R}_{>0}.$$

In this case (11) transforms into:

$$r \frac{1}{2\sqrt{V}}(-kV + c\sqrt{V}r\sqrt{V}) + \frac{\gamma}{\sqrt{p}}\sqrt{V} = \frac{\sqrt{V}}{2} \left(-kr + cr^2 + \frac{2\gamma}{\sqrt{p}} \right) \leq 0, \quad (35)$$

and choosing

$$\gamma < \frac{\sqrt{p}}{2} (kr - cr^2) \quad (36)$$

ensures that (11) is satisfied for all V . Optimizing the right-hand side of (36) for r we obtain that the maximum is achieved for $r = k/(2c)$. Hence, the condition

$$\gamma < \frac{1}{8} \frac{\sqrt{p}k^2}{c} = \frac{1}{16} \frac{pk^2}{\sigma B}.$$

ensures that (35) holds, and that the solutions of (30) passing through

$$\Omega_a^* = \{(x, \lambda) \mid x \in \mathbb{R}^n, \lambda \in \mathbb{R}, \lambda(t_0) \geq \sqrt{V(x(t_0))}k\sqrt{p}/(4\sigma B)\}$$

are bounded and remain in Ω_a^* . Given that $\sqrt{p}\|x(t_0)\| \leq \sqrt{V} \leq \sqrt{p}\|x(t_0)\|$ we can conclude that

$$\Omega'_a = \{(x, \lambda) \mid x \in \mathbb{R}^n, \lambda \in \mathbb{R}, \lambda(t_0) \geq \|x(t_0)\|k\sqrt{pp}/(4\sigma B)\} \subset \Omega_a^*$$

and

$$\Omega_a = \{(x, \lambda) \mid x \in \mathbb{R}^n, \lambda \in \mathbb{R}, \lambda(t_0) \geq \|x(t_0)\|kp/(4\sigma B)\} \supset \Omega_a^*$$

Hence solutions starting in Ω'_a will remain in Ω_a . $\square$

We illustrate with the next example, the bounds for γ established in Corollary 3 can be substantially less conservative than those provided by the small-gain theorem [22] for the same system.

Example 6 Consider the following cascaded system

$$\begin{aligned} \dot{x}_1 &= -\tau_1 x_1 + c_1 x_2 \\ \dot{x}_2 &= -c_2 |x_1|, \quad \tau_1, c_1, c_2 \in \mathbb{R}_{>0}. \end{aligned} \quad (37)$$

According to Corollary 4.1 from [22] the following choice of system parameters ensures existence of initial conditions corresponding to nontrivial bounded (in forward time) solutions:

$$0 < c_2 < \frac{1}{16} \frac{\tau_1^2}{c_1}. \quad (38)$$

Now we will use Corollary 3 to estimate the bounds for c_2 that guarantee existence of nontrivial bounded solutions. As a Lyapunov candidate we chose $V(x_1) = x_1^2$. Hence

$k = -2\tau_1$, $\sigma = 1$, $B = c_1$, $\bar{p} = \underline{p} = 1$. Then, according to Corollary 3 the following inequality guarantees existence of nontrivial bounded solutions for (37):

$$0 < c_2 < \frac{4\tau_1^2}{16} \frac{1}{c_1} = \frac{1}{4} \frac{\tau_1^2}{c_1}.$$

The domain of admissible values for c_2 that follows from Corollary 3 is four times larger than that specified by (38) which is based on the non-uniform small-gain theorem. This illustrates that our present result is less conservative for the problems of this class.

4 Discussion and Conclusions

In this brief manuscript we provided further results that allow to address the issue of positive invariance and convergence for a range of problems in control and in the broader domain of dynamical systems. In particular, we considered systems in which stable motions in higher dimensional subspace of the system state space are coupled with unstable motions in a lower dimensional subspace. The main motivation is to go beyond current restrictions of convergence results in [22], [4] and provide an analysis method for a relevant class of systems with unstable solutions.

The results we provided do not require (partial) knowledge of the system solution; they do not assume availability of input-output gains and estimates of contraction rates for the stable parts of the system. The advantage of our method is that it has the flavor of simplicity inherent to conventional Lyapunov-based analysis. Furthermore, the boundedness/unboundedness criteria have similar shape with that of the celebrated Lyapunov stability/instability conditions.

Our method allows to produce substantially less conservative convergence results than other available techniques such as [22], [4]. We illustrated this possibility with a simple example. Availability of these less-conservative estimates is advantageous in parameter estimation problems [21], [20] for in these applications the upper bound for γ determines the maximal speed of exploration in the space of parameters. The higher the speed, the better the overall performance.

In addition to the boundedness criteria, geometric intuition behind our results allows to estimate the domains of initial conditions corresponding to solutions which necessarily escape a given small neighborhood of the origin. These results can be applied to the problem of estimating relaxation times in dynamical systems [3].

References

- [1] G. Bastin and M. Gevers. Stable adaptive observers for nonlinear time-varying systems. *IEEE Trans. on Automatic Control*, 33(7):650–658, 1988.
- [2] C. I. Byrnes, F.D. Priscoli, A. Isidori, and W Kang. Structurally stable output regulation of nonlinear systems. *Automatica*, 33(3):369–385, 1997.
- [3] A.N. Gorban. Singularities of transition processes in dynamical systems: Qualitative theory of critical delays electron. *Electr. J. Diff. Eqns. Monograph* 5, 2004. <http://ejde.math.txstate.edu/Monographs/05/>.

- [4] A. Ilchman. Universal adaptive stabilization of nonlinear systems. *Dynamics and Control*, (7):199–213, 1997.
- [5] A. Isidori. *Nonlinear control systems: An Introduction*. Springer-Verlag, second edition, 1989.
- [6] L. Ljung. *System Identification: Theory for the User*. Prentice Hall, 1999.
- [7] A.M. Lyapunov. *The general problem of the stability of motion (in Russian)*. Kharkov Mathematical Society, 1892. Republished by the University of Toulouse, 1908 and Princeton University Press, 1949 (in French). Republished in English by Int. Journal of Control, 1992.
- [8] R. Marino. Adaptive observers for single output nonlinear systems. *IEEE Trans. Automatic Control*, 35(9):1054–1058, 1990.
- [9] R. Marino and P. Tomei. Global adaptive output-feedback control of nonlinear systems, part I: Linear parameterization. *IEEE Trans. Automatic Control*, 38(1):17–32, 1993.
- [10] I. Miroshnik, V. Nikiforov, and A. Fradkov. *Nonlinear and Adaptive Control of Complex Systems*. Kluwer, 1999.
- [11] A.S. Morse. Control using logic-based switching. In *Trends in Control*, pages 69–113. Springer-Verlag, 1995.
- [12] K. S. Narendra and A. M. Annaswamy. *Stable Adaptive systems*. Prentice-Hall, 1989.
- [13] H. Nijmeijer and A. van der Schaft. *Nonlinear Dynamical Control Systems*. Springer-Verlag, 1990.
- [14] Jean-Baptiste Pomet. Remarks on sufficient information for adaptive nonlinear regulation. In *Proceedings of the 31-st IEEE Control and Decision Conference*, pages 1737–1739, 1992.
- [15] M. I. Rabinovic, R. Huerta, P. Varona, and V.S. Aframovich. Transient cognitive dynamics, metastability, and decision-making. *PLOS Computational Biology*, 4(5):1–9, 2008.
- [16] M. I. Rabinovic, P. Varona, A.I. Selverston, and H.D.I. Abarbanel. Dynamical principles in neuroscience. *Reviews of Modern Physics*, 78:1213–1265, 2006.
- [17] R. Rockafellar. *Convex Analysis*. Princeton University Press, 1970.
- [18] E. Steur, I.Yu. Tyukin, H. Nijmeijer, and C. van Leeuwen. Reconstructing dynamics of spiking neurons from input-output measurements in vitro. In *Proceedings of the 3-d IEEE Physics and Control Conference*. Potsdam, Germany, 3 – 7 September 2007.
- [19] I.Yu. Tyukin, Prokhorov, and C. van Leeuwen. Adaptation and parameter estimation in systems with unstable target dynamics and nonlinear parametrization. *IEEE Trans. on Automatic Control*, 52(9):1543–1559, 2007.

- [20] I.Yu. Tyukin, D. V. Prokhorov, and C. van Leeuwen. Adaptive classification of temporal signals in fixed-weights recurrent neural networks: an existence proof. *Neural Computation*, 20(10):2564–2596, 2008.
- [21] I.Yu. Tyukin, E. Steur, H. Nijmeijer, and C. van Leeuwen. Adaptive observers and parametric identification for systems in non-canonical adaptive observer form. *Submitted*.
- [22] I.Yu. Tyukin, E. Steur, H. Nijmeijer, and C. van Leeuwen. Non-uniform small-gain theorems for systems with unstable invariant sets. *SIAM Journal on Control and Optimization*, 47(2):849–882, 2008.
- [23] I.Yu. Tyukin, E. Steur, H. Nijmeijer, and C. van Leeuwen. State and parameter estimation for systems in non-canonical adaptive observer form. In *Proceedings of the 17-th IFAC World Congress on Automatic Control*, 2008.
- [24] V.I. Zubov. *Methods of A.M. Lyapunov and Their Applications*. P. Noordhoff, Ltd., Groningen, 1964.

5 Appendix. Star-shaped sets and envelopes

Convex analysis is a well known and widely used subject in the domain of optimization [17]. In contrast, the idea of star shaped functions is not so well known. Here we provide a list of definitions and properties related to the notions of convexity and star-shapedness that are referred to in the main text of the manuscript.

Let V be a real vector space. A closed interval $[x, y]$ is defined as a segment connecting two vectors $x, y \in V$: $[x, y] = \{\gamma x + (1 - \gamma)y \mid \gamma \in [0, 1]\}$.

Definition 1 *A set $S \subset V$ is star shaped with respect to a point $x \in S$, if for any $y \in S$ the interval $[x, y]$ also belongs to S : $[x, y] \subset S$.*

The following properties hold for the star shaped sets in V :

- A set is convex iff it is star shaped with respect to its every point.
- Let $x \in V$ and W be a family of star shaped sets with respect to x . Then both the intersection $\cap_{S \in W} S$ of sets from W and the union $\cup_{S \in W} S$ of sets from W are star shaped with respect to x .
- Let E be a real vector space, $A : V \rightarrow E$ be a linear map, and $S \subset V$ be a star shaped set with respect to a point $x \in S$. Then the image $A(S)$ of A is star shaped with respect to $A(x)$.

Definition 2 *For any set $D \subset V$ and a point $x \in D$ the star shaped envelope of D with respect to x , $\text{star}_x(D)$, is the minimal star shaped set with respect to x which includes D .*

Star shaped envelop exists and could be defined in two alternative ways. Namely, “from above” (as an intersection):

$$\text{star}_x(D) = \bigcap_{S \in W_x(D)} S,$$

where $W_x(D)$ is a family of sets that are star shaped with respect to x and include D , and “from below” (as union of intervals):

$$S = \bigcup_{y \in D} [x, y]. \quad (39)$$

Notice that deriving a star shaped (w.r.t. a point) envelope of a set is a much easier procedure computationally than that of deriving a convex envelop of the same set.

Let us remind that epigraph of a real valued function $f : S \rightarrow \mathbb{R}$ is a subset of $S \times \mathbb{R}$ that consists of all points lying on or above its graph:

$$\text{epi}(f) = \{(x, \gamma) \mid x \in S, \gamma \geq f(x)\}.$$

Definition 3 A function $f : S \rightarrow \mathbb{R}$ is star shaped with respect to $x \in S$, if $\text{epi}(f)$ is star shaped set with respect to $(x, f(x))$.

Clearly, S must necessarily be star shaped with respect to x . Alternatively, we can use the following definition.

Definition 4 A function $f : S \rightarrow \mathbb{R}$ (on a star shaped set S with respect to x) is star shaped with respect to x iff the following holds for any $y \in S$ and every $\gamma \in [0, 1]$:

$$f(\gamma x + (1 - \gamma)y) \geq \gamma f(x) + (1 - \gamma)f(y).$$

This is a form of Jensen’s inequality with one “fixed end”. It is obvious that a function on S is convex iff it is star shaped with respect to every point $x \in S$.

Definition 5 Let $f : S \rightarrow \mathbb{R}$ be a bounded from above and below function ($A < f(x) < B$). Supremum of star-shaped (with respect to x) minorants of f is the star shaped envelope of f with respect to x , $\text{star}_x(f)$.

Let $S \subset V$ be a star shaped set with respect to $x \in S$. The following list of properties holds for the star shaped sets and functions:

1.

$$\text{epi}(\text{star}_x(f)) = \text{star}_x(\text{epi}(f)). \quad (40)$$

2.

$$\text{conv}(f)(y) \leq \text{star}_x(f)(y) \leq f(y). \quad (41)$$

The first property follows immediately from the definition, and the second property is a consequence of the following fact:

$$\text{epi}(f) \subset \text{epi}(\text{star}_x(f)) \subset \text{epi}(\text{conv}_x(f)). \quad (42)$$

From the definition of a star envelope of a set from below (the last expression in (39)) we can produce the definition of a star envelope of a function. Let us start from 1D case (i.e. $V = \mathbb{R}$). Let $S = [a, b]$, $x \in S$. For any $z \in S$ ($z \neq x$) we define

$$\phi_{z,x}(y) = \begin{cases} \min \left\{ f(y), \frac{y-x}{z-x}f(x) + \frac{z-y}{z-x}f(z) \right\} & \text{if } 0 < \frac{y-x}{z-x} < 1; \\ f(y) & \text{else.} \end{cases} \quad (43)$$

As follows from the Jensen inequality with one fixed end, the following property holds

$$\text{star}_x(f)(y) = \inf_{z \in S} \{\phi_{z,x}(y)\}$$

The following proposition is obvious:

Proposition 1 *Let $f : S \rightarrow \mathbb{R}$ be a continuous function. Then the following properties hold:*

- 1) *The functions $\phi_{z,p}(y)$, $p \in S$ are equicontinuous with f ;*
- 2) *The set of functions $\{\phi_{z,p}(y)\}$, $p \in S$ is compact;*
- 3) *The function $\text{star}_p(f)(y)$, $p \in S$ exists and is a continuous function.*
- 4) *If a continuous function on $[a, b]$ achieves its minimum at a single point x , then its star shaped envelope with respect to x has the same property.*
- 5) *The star shaped envelope (w.r.t. a point $p \in S$) of a monotone function is monotone.*

The first property follows immediately from (43). Properties 2,3 follow from the Arzela–Ascoli theorem. Indeed, $\{\phi_{z,p}(y)\}$, $p \in S$ is pointwise relatively compact and closed. Hence, taking property 1 into account, we can conclude that the set of functions $\{\phi_{z,p}(y)\}$ is compact. To demonstrate existence and continuity of $\text{star}_p(f)(y)$, $p \in S$ consider a sequence $\{g_i\}_{i=1}^\infty$ of grids $g_i = \{a, a + (b-a)/i, a + (b-a)k/i, \dots, b\}$, $1 \leq k < i$ on S , and define

$$f_i(y) = \inf_{r \in g_{2i}} \{\phi_{r,p}(y)\}$$

The sequence of functions $\{f_i(y)\}_{i=1}^\infty$ is equicontinuous and $f_1 \geq f_2 \geq \dots \geq f_n \geq$ pointwise. This means that the sequence converges uniformly, and that $\lim_{i \rightarrow \infty} f_i(y) = \inf_{z \in S} \{\phi_{z,p}(y)\} = \text{star}_p(f)(y)$. Property 4 is easily verifiable by the contradiction argument. Property 5 is the consequence of that functions $f_i(y)$ are monotone by construction and are equicontinuous. $\square$

Let $V = \mathbb{R}^n$, and $S \subset V$ be a compact and star shaped with respect to x . For every $z \in S$, and $\gamma \in [0, 1]$ we define

$$\psi_z(\gamma) = \min\{f((1-\gamma)x + \gamma z), (1-\gamma)f(x) + \gamma f(z)\}. \quad (44)$$

Then

$$\text{star}_x(f)(y) = \inf_{(1-\gamma)x + \gamma z = y, \gamma \in [0, 1], z \in S} \{\psi_z(y)\}.$$

The properties of function $\text{star}_x(f)(y)$ depend on the properties of mapping

$$y \mapsto \{(\gamma, z) \mid (1-\gamma)x + \gamma z = y, \gamma \in [0, 1], z \in V\}. \quad (45)$$

If this mapping is continuous (in the Hausdorff metrics in the space of compact sets), then the star shaped envelope of every continuous in S function with respect to x is also continuous. Mapping (45) is continuous in S iff the Minkovski functional

$$p(y) = \inf_{a > 0} \{a \mid y - x \in a(S - x)\}.$$

is continuous in S .